

THE CLOSEST DAMPED LYMAN ALPHA SYSTEM

ERIC D. MILLER

Dept. of Astronomy, University of Michigan, Ann Arbor, MI 48109-1090 emiller@astro.lsa.umich.edu

PATRICIA M. KNEZEK

Dept. of Physics & Astronomy, The Johns Hopkins University, Baltimore, MD 21218-2695 pmk@pha.jhu.edu

JOEL N. BREGMAN

Dept. of Astronomy, University of Michigan, Ann Arbor, MI 48109-1090 jbregman@umich.edu

Accepted by Astrophysical Journal Letters

ABSTRACT

A difficulty of studying damped Ly α systems is that they are distant, so one knows little about the interstellar medium of the galaxy. Here we report upon a damped Ly α system in the nearby galaxy NGC 4203, which is so close ($v_{helio} = 1117 \text{ km s}^{-1}$) and bright ($B_o = 11.62$) that its H I disk has been mapped. The absorption lines are detected against Ton 1480, which lies only $1.9'$ ($12 h_{50}$ kpc) from the center of NGC 4203. Observations were obtained with the Faint Object Spectrograph on HST (G270H grating) over the $2222\text{--}3277 \text{ \AA}$ region with 200 km s^{-1} resolution. Low ionization lines of Fe, Mn, and Mg were detected, leading to metallicities of -2.29 , < -0.68 , and > -2.4 , which are typical of other damped Ly α systems, but well below the stellar metallicity of this type of galaxy. Most notably, the velocity of the lines is $1160 \pm 10 \text{ km s}^{-1}$, which is identical to the H I rotational velocity of 1170 km s^{-1} at that location in NGC 4203, supporting the view that these absorption line systems can be associated with the rotating disks of galaxies. In addition, the line widths of the Mg lines give an upper limit to the velocity dispersion of 167 km s^{-1} to the 99% confidence level.

Subject headings: galaxies: abundances — galaxies: individual (NGC 4203) — galaxies: ISM — quasars: absorption lines

1. INTRODUCTION

Damped Ly α systems (hereafter, DLA) are the primary tracers of metallicity over cosmological distances (*e.g.*, Lu *et al.* 1996) and provide a method of identifying distant galaxies (*e.g.*, Giallisco, Steidel, and Szalay 1994). It is likely that these DLA are associated with galaxies, but whether the lines arise in a rotating disk or in the halo remains a matter of controversy (Charlton and Churchill 1996; Prochaska and Wolfe 1997; Lu, Sargent, and Barlow 1997). One reason for this uncertainty is that we possess little information on the cold gas content of these galaxies. H I velocity maps of the absorbing hosts would be very valuable in addressing this issue, but even the two previously known nearest DLAs, at $z = 0.101$ and $z = 0.167$ (Petitjean *et al.* 1996; Lanzetta *et al.* 1997), are too distant for such observations to be feasible.

Another issue is the meaning of the measured abundances, which nominally indicate that the metallicities of DLA are typically 10–300 times below the solar value over redshifts of $z = 0\text{--}3$. (For higher redshifts, the metallicity appears to be even lower; Lu, Sargent, and Barlow 1998.) However, low apparent abundances are also derived for sight lines out of the Milky Way, but in this case we know this is due to depletion onto grains (*e.g.*, Cardelli, Sembach, and Savage 1995; Savage and Sembach 1996). Unfortunately, depletion onto grains can raise a substantial uncertainty in the metallicity deduced from DLAs.

Some insight into these issues is offered by our observations of a DLA caused by a galaxy so nearby that some of the properties of its H I disk are already known. Here we present ultraviolet resonance line observations that led to the discovery of a DLA system toward the quasar Ton 1480

($z = 0.614$; $V = 16.5$), which lies behind the nearby galaxy NGC 4203 at a projected separation of only $1.9'$ ($12 h_{50}$ kpc) from the center. NGC 4203 is a nearby ($v_{helio} = 1117 \text{ km s}^{-1}$), bright ($B_o = 11.62$) early-type galaxy that is not in a cluster, has normal colors, with $D_{25} = 3.6'$, and a one-dimensional velocity dispersion of 175 km s^{-1} . The background quasar, Ton 1480, was discovered by us as part of a program whereby we use background X-ray sources to identify quasars near targets of interest, such as galaxies and galaxy clusters (Knezek and Bregman 1998). This galaxy has been observed in many wavebands, including an H I map (van Driel *et al.* 1988) with both column density and velocity information for the neutral gas. In the following, we discuss the absorption line observations, the column densities for Fe, Mg, and Mn, the kinematics of the absorbing gas, and how this relates to the known kinematics of the H I gas.

2. DATA

The data were taken in January 1997 with the Faint Object Spectrograph aboard HST. The 1640 second exposure was taken through the $0.86''$ aperture using the G270H grating and Red Digicon detector. In this configuration, the expected FOS instrumental line width (FWHM) is $1.97 \text{ \AA}$, providing a spectral resolution of 211 km s^{-1} at the Mg II $\lambda 2800$ doublet. The wavelength coverage ranges from 2222 to $3277 \text{ \AA}$, with a S/N ratio of about 20 throughout the spectrum.

Visual inspection of the spectrum reveals several strong absorption features (see Figure 1). The Mg II $\lambda 2800$ doublet from NGC 4203 can be seen shifted $10.8 \text{ \AA}$ (1160 km s^{-1}) from its Galactic counterpart. The Fe II $\lambda 2600$ line

also can be seen shifted $10.1 \text{ \AA}$ (1165 km s^{-1}) from the Galactic line. In general, lines with $W_\lambda > 0.5 \text{ \AA}$ are easily visible to the eye. Other such lines include Fe II $\lambda 2374$, $\lambda 2382$, as well as their corresponding Galactic lines.

To detect fainter absorption features, we utilized the line-searching software developed for the HST absorption line key project (Schneider *et al.* 1993). This algorithm performs both Gaussian and PSF fitting to absorption features across the spectrum, finding line centers and equivalent widths with errors. The detection threshold was set to $3\text{-}\sigma$, which corresponds to a minimum equivalent width of $0.15 \text{ \AA}$ across most of the spectrum. The fitting procedure detected 20 lines (see Table 1); of these, 7 are Galactic in origin, 6 are from the DLA, 3 are blends of Milky Way and DLA lines, and 4 are unidentified. For a spectrum with 2064 resolution elements, we would expect roughly 1 spurious detection above the $3\text{-}\sigma$ limit. Thus a few of the unknown lines, which are all near the $3\text{-}\sigma$ limit, could possibly be real lines; however, they are inconsistent with any known atomic resonance lines, either Galactic or at the velocity of NGC 4203. It remains to be seen if they are real absorption lines from intervening material between the quasar and NGC 4203.

We identified weak lines in Fe II, Mg I, and Mn II at the redshift of NGC 4203, as well as the easily visible lines mentioned above. All identified DLA lines are listed in Table 1 which gives the vacuum rest wavelength, line center, and equivalent width with $1\text{-}\sigma$ errors.

Analysis of the Galactic features sheds some light on the quality of the HST/FOS wavelength calibration. At the high Galactic latitude of Ton 1480 ($l = 173^\circ$, $b = 80.1^\circ$), the Galactic ISM should be nearly at rest relative to the heliocentric velocity. Thus any Galactic line should be barely shifted from its rest wavelength in the direction of Ton 1480.

To determine the average expected shift in the Milky Way lines, we calculated a weighted mean line-of-sight velocity for the H I disk,

$$\langle v_{\text{LOS}} \rangle = \frac{\iint v_{\text{LOS}}(R, z) n(R, z) dR dz}{\iint n(R, z) dR dz},$$

integrating out to $z = 500 \text{ pc}$. Using the estimate of the H I density z -dependence from Dickey and Lockman (1990) and the standard equation for differential Galactic rotation, $v_{\text{LOS}} = \Theta_\odot [R_\odot/R - 1] \sin l \cos b$, we find $\langle v_{\text{LOS}} \rangle = -4 \text{ km s}^{-1}$, only 2% of our spectral resolution.

From the FOS calibrated (vacuum) wavelengths, the average shift of the clean Galactic features is $-1.23 \pm 0.2 \text{ \AA}$. This corresponds to a shift of $-130 \pm 20 \text{ km s}^{-1}$ at $2800 \text{ \AA}$, much larger than the expected shift found above. Thus the FOS wavelength calibration is off by about $1.2 \text{ \AA}$ to the blue. This is within the pointing error for a FOS Binary Target Acquisition; in addition, it has been found recently that the zero-point of the FOS wavelength scale varied considerably over the lifetime of the instrument (Keyes 1998). Since we did not obtain a lamp spectrum at the time of the observation but instead relied on the default calibration used in the HST pipeline, our wavelength calibration could indeed be off by this amount.




FIG. 1.— Selected regions of the normalized spectrum of Ton 1480. The labels correspond to ID entries of absorption lines in Table 1.

Consequently, we used the Galactic lines (plus the 4 km s^{-1} offset) as the fiducial zero velocity. The velocity shifts in all DLA lines from the Galactic zero-point are given in Table 1. (For display purposes, all wavelength measurements have been shifted $1.2 \text{ \AA}$ to the red.) The average line shift for the unblended lines is $1163 \pm 10 \text{ km s}^{-1}$ from the Galactic reference lines, implying a heliocentric velocity of $v_{\text{helio}} = 1159 \pm 10 \text{ km s}^{-1}$.

As mentioned earlier, several absorption lines are blended with Milky Way features. The Fe II $\lambda 2374$, Fe II $\lambda 2587$ and Mn II $\lambda 2577$ lines are all blended with Galactic Fe II lines. This Mn II line is the only line of this element detected, so only an upper limit can be obtained for its abundance. For Fe, three lines (Fe II $\lambda 2344$, Fe II $\lambda 2383$, and Fe II $\lambda 2600$) were cleanly detected. Additionally, non-detections of two Fe II lines and three Fe I lines provide upper limits for their equivalent widths. Finally, we have detected three Mg lines in two stages of ionization, the Mg II $\lambda 2800$ doublet and Mg I $\lambda 2853$. The doublet is clean and pronounced while the Mg I line is weak and appears broadened. It also has an anomalous velocity, shifted 75 km s^{-1} from the mean determined above. It is possible this line is blended, although there are no strong Galactic features in this region of the spectrum. Thus its nature remains questionable, and we treat it only as an upper limit.

3. INTERPRETATION & CONCLUSIONS

Column densities can be determined for weak (optically thin) lines by using the relation (*e.g.*, Spitzer 1978):

$$N = 1.13 \times 10^{20} W_\lambda (\text{\AA}) / f \lambda^2 (\text{\AA}) \text{ cm}^{-2}$$

The unblended Fe II lines are all thin lines, since the value of W_λ / λ^2 scales linearly with the oscillator strength, f (taken from Morton 1991). The Mg II $\lambda 2800$ doublet lines appear to be on the Doppler portion of the curve of growth, since the line strength does not scale as the oscillator strength. Since we have two lines, we can simply take the line ratio, and we find that $W_{\lambda 2803} / W_{\lambda 2796} = 0.71 \pm 0.05$.

TABLE 1
NGC 4203 ABSORPTION LINE DATA

Species	ID ^a	$\lambda(\text{\AA})$	$\lambda_{obs}(\text{\AA})$	$v_r(\text{km/s})$	$W_\lambda(\text{\AA})$	$N(\text{cm}^{-2})$
Fe I	—	2484.02	2494.5	—	≤ 0.091	$\leq 3.0 \times 10^{12}$
Fe I	—	2523.61	2533.3	—	≤ 0.082	$\leq 5.2 \times 10^{12}$
Fe I	—	2719.83	2730.4	—	≤ 0.104	$\leq 1.3 \times 10^{13}$
Fe II	2	2344.21	2353.5 ± 0.4	1168.9	0.280 ± 0.064	$5.3 \pm 1.2 \times 10^{13}$
Fe II	5	2382.77	2392.2 ± 0.1	1124.8 ^b	0.804 ± 0.049	$5.3 \pm 0.3 \times 10^{13}$
Fe II	11	2600.17	2610.3 ± 0.1	1164.5	0.683 ± 0.041	$5.1 \pm 0.3 \times 10^{13}$
Fe II	—	2249.88	2259.0	—	≤ 0.129	$\leq 1.1 \times 10^{15}$
Fe II	—	2260.78	2270.7	—	≤ 0.127	$\leq 7.6 \times 10^{14}$
Fe II	4	2374.46	2383.2 ± 0.1	1117.4 ^c	$\leq 0.773^c$	$\leq 5.5 \times 10^{14}$
Fe II	9	2586.65	2597.3 ± 0.2	1225.1 ^{b,c}	$\leq 0.281^c$	$\leq 7.4 \times 10^{13}$
Mg I	17	2852.96	2864.6 ± 0.3	1238.9	0.203 ± 0.052	$\leq 1.5 \times 10^{12d}$
Mg II	14	2796.35	2807.1 ± 0.1	1156.8	1.210 ± 0.039	$\geq 2.9 \times 10^{13}$
Mg II	15	2803.53	2814.5 ± 0.1	1162.4	0.857 ± 0.042	$\geq 4.0 \times 10^{13}$
Mn II	8	2576.88	2586.7 ± 0.1	1180.8 ^c	$\leq 0.649^c$	$\leq 3.1 \times 10^{13}$
Mn II	—	2594.50	2604.6	—	≤ 0.097	$\leq 6.0 \times 10^{12}$
Mn II	—	2606.46	2616.7	—	≤ 0.098	$\leq 8.5 \times 10^{12}$
Gal Fe II	1	2344.21	2344.4 ± 0.2	—	0.493 ± 0.056	—
Gal Fe II	3	2374.46	2374.4 ± 0.3	—	0.258 ± 0.059	—
Gal Mn II	7	2576.88	2576.6 ± 0.2	—	0.270 ± 0.044	—
Gal Fe II	10	2600.17	2600.2 ± 0.1	—	0.675 ± 0.041	—
Gal Mg II	12	2796.35	2796.3 ± 0.1	—	1.000 ± 0.063	—
Gal Mg II	13	2803.53	2803.7 ± 0.1	—	1.060 ± 0.072	—
Gal Mg I	16	2852.96	2852.8 ± 0.3	—	0.224 ± 0.051	—
unknown	6	—	2477.0 ± 0.4	—	0.140 ± 0.045	—
unknown	18	—	3029.6 ± 0.4	—	0.179 ± 0.050	—
unknown	19	—	3089.8 ± 0.4	—	0.287 ± 0.095	—
unknown	20	—	3273.0 ± 0.4	—	0.250 ± 0.084	—

^aAbsence of a line ID indicates a non-detection.

^bThe Galactic counterparts of these lines are blended, so it is difficult to determine an accurate velocity.

^cThese DLA lines are blended, leading to inaccurate velocities and upper limits on W_λ and N .

^dThe anomalous velocity of the Mg I line makes its nature questionable, thus we include it as an upper limit.

In the thin-line limit, we would expect the line ratio to be $W_{\lambda 2803}/W_{\lambda 2796} = (f\lambda^2)_{\lambda 2803}/(f\lambda^2)_{\lambda 2796} = 0.50$, so these lines are indeed optically thick, and we can only obtain a lower limit to the Mg II column density. The Mg I and Mn II lines are probably thin, but it is problematic to determine column densities without more lines from these species.

Column densities and 1- σ errors for each line are given in Table 1. For lines which were not detected, a 3- σ upper limit to the equivalent width can be determined from consideration of the detection threshold and noise. Assuming the lines are weak, these values represent an upper limit to the column density.

To obtain the abundance of an element relative to solar, an accurate measurement of the H I column is needed. For this system, there are several possible methods. Van Driel *et al.* (1988) used 21-cm observations, finding an H I column of about $1 \times 10^{20} \text{ cm}^{-2}$ towards Ton 1480. A more accurate estimate comes from X-ray absorption observations. The X-ray absorption column toward Ton 1480 was determined for the *ROSAT PSPC* data, which yielded a value of $3.38 \times 10^{20} \text{ cm}^{-2}$ (Bregman, Hogg, and Roberts

1995), while the Galactic H I column in this direction is only $1.14 \times 10^{20} \text{ cm}^{-2}$ (Hartmann and Burton 1997). Since the value of the Galactic H I column is the same as the Galactic X-ray absorption column (Arabadjis and Bregman 1998, and references therein), we attribute the excess absorption column to that of the disk of NGC 4203, which leads to a value of $N_{\text{H}} = 2.2 \times 10^{20} \text{ cm}^{-2}$. To further constrain this value, Ly α observations are planned for HST in the near future.

To obtain the total elemental abundance, we need to know the column density in each stage of ionization. We are severely limited since we have detected only one ionization stage for two of the three elements. Thus we need to model the ionization structure to correct for this. The ionization model in this region of NGC 4203 was produced using CLOUDY (Ferland *et al.* 1998). Two different ionizing spectra (Ikeuchi and Ostriker 1986; Haardt and Madau 1996) were tested, along with the two values for the H I column N_{H} given above. For most of our lines, there is little difference in the ionization among the four models. Fe I and Fe II account for $\sim 5\%$ and $\sim 90\%$ of the Fe in each model, respectively. The fractional ionization of

Mg is Mg I ~ 10 –20% and Mg II ~ 70 –75%. It should be noted that adjusting N_{H} by a factor of 2 makes less of a difference than changing the spectrum of the ionizing radiation. This is especially true for Fe I, where the difference between the two model spectra is a factor of 2 in the fractional ionization.

We used the spectrum from Figure 5a of Haardt and Madau (1996), normalized to $J_{\nu}(912 \text{ \AA}) = 2 \times 10^{-23} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$ (Dove and Shull 1994), and the X-ray column of $N_{\text{H}} = 2.2 \times 10^{20} \text{ cm}^{-2}$. The model gives the following ionization fractions: Fe I = 0.038, Fe II = 0.942, Mg I = 0.117, Mg II = 0.764, and Mn II = 0.635. The relative column densities for Fe I and Fe II are consistent with this model. However, according to our data $N(\text{Mg}^0)/N(\text{Mg}^+) = 0.04$, whereas our model predicts it to be 0.15. It appears the ionization in this region is higher than our model predicts; this is especially likely given the anomalous velocity of the Mg I line, which could indicate that the Mg I and Mg II absorbers are not co-spatial.

Earlier results indicate that ionization correction factors for Mg II and Fe II may be insignificant for this observed H I column density (Viegas 1995; Prochaska and Wolfe 1996). We find our data to be consistent with these results, and so we will not apply any ionization correction to these species for the following analysis.

The column densities listed in Table 1 give the following results for the heavy element abundances, relative to the solar values of $\log(\text{Fe}) = 7.67$, $\log(\text{Mn}) = 5.39$, $\log(\text{Mg}) = 7.58$, and $\log(\text{H}) = 12.00$ (Anders and Grevesse 1989):

$$[\text{Fe}/\text{H}] = -2.29 \pm 0.10$$

$$[\text{Mn}/\text{H}] < -0.68$$

$$[\text{Mg}/\text{H}] > -2.4$$

These values are significantly below the expected stellar metallicity, although it is difficult to separate true under-abundances from depletion onto grains since each of these

elements have high condensation temperatures.

A comparison between the known velocity of the H I disk with the absorption line velocities can help us understand the nature of the absorption site. Van Driel's (1988) velocity maps give an H I velocity of 1170 km s^{-1} along this line of sight in NGC 4203. From the centers of the unblended Fe II and Mg II absorption lines, the average heliocentric velocity is found to be $1160 \pm 10 \text{ km s}^{-1}$. This is indistinguishable from the H I value, clearly demonstrating that the absorption is due to a galactic disk of gas.

Limits can also be placed on the internal motions of the absorbing gas. The widths of the strongest Fe II lines average to $260 \pm 27 \text{ km s}^{-1}$ FWHM, with an instrumental line width of 241 km s^{-1} , which sets an upper limit to the velocity dispersion of 224 km s^{-1} at the 99% confidence level. For Mg II, the constraint is stricter, with an average width of $248 \pm 12 \text{ km s}^{-1}$ FWHM and an instrumental profile of 210 km s^{-1} yielding a maximum velocity dispersion of 167 km s^{-1} at 99% confidence.

In the future, we hope to use this unique system to pursue additional issues of DLAs. A planned HST observation with STIS should determine whether high ionization state gas is present, permit column densities to be measured for other species, and provide a third measure of the H I column. Higher spectral resolution studies will be able to examine the velocity structure of the absorbing material. Finally, the intrinsic metallicity of the system can be studied through optical observations of H II regions and by X-ray studies.

We would like to thank J. Bergeron, J. Charlton, D. Schneider, C. Cowley, E. Schulman, and the staff at STScI for their advice and assistance. Financial support for this work has been provided by NASA through NAG5-3247 and STScI GO-06547.

REFERENCES

- Anders, E. and Grevesse, N. 1989, *Geo. Cosmo. Acta*, 53, 197
 Arabadjis, J. S., and Bregman, J. N. 1998, *ApJ*, in press
 Bregman, J. N., Hogg, D. E., and Roberts, M. S. 1995, *ApJ*, 441, 561
 Cardelli, J. A., Sembach, K. R. and Savage, B. D. 1995, *ApJ*, 440, 241
 Charlton, J. C., and Churchill, C. W. 1996, *ApJ*, 465, 631
 Dickey, J. M., and Lockman, F. J. 1990, *ARAA*, 28, 215
 Dove, J. B., and Shull, J. M. 1994, *ApJ*, 423, 196
 Ferland, G. J., Korista, K. T., Verner, D. A., Ferguson, J. W., Kingdon, J. B., and Verner, E. M. 1998, *PASP*, 110, 761
 Gialavisco, M., Steidel, C. C., and Szalay, A. S. 1994, *ApJ*, 425, L5
 Haardt, F., and Madau, P. 1996, *ApJ*, 461, 20
 Hartmann, D., and Burton, W. B. 1997, *Atlas of Galactic Neutral Hydrogen*, (Cambridge: Cambridge University Press)
 Ikeuchi, S., and Ostriker, J. P. 1986, *ApJ*, 301, 522
 Keyes, C. D., ed. 1998, *HST Data Handbook*, Version 3.1, Volume II (Baltimore: STScI)
 Knezek, P. M., and Bregman, J. N. 1998, *AJ*, 115, 1737
 Lanzetta, K. M. et al. 1997, *AJ*, 114, 1337
 Lu, L., Sargent, W. L. W., and Barlow, T. A. 1997, *ApJ*, 484, 131
 Lu, L., Sargent, W. L. W., and Barlow, T. A. 1998, to appear in *Cosmic Chemical Evolution*, (Kluwer: Dordrecht), also *astro-ph/9710370*
 Lu, L., Sargent, W. L. W., Barlow, T. A. Churchill, C., and Vogt, S. S. 1996, *ApJS*, 107, 475
 Morton, D. C. 1991, *ApJS*, 77, 119
 Petitjean, P., Theodore, B., Smette, A., and Lespine, Y. 1996, *A&A*, 313, L25
 Prochaska, J. X., and Wolfe, A. M. 1996, *ApJ*, 470, 403
 Prochaska, J. X., and Wolfe, A. M. 1997, *ApJ*, 487, 73
 Savage, B. D., and Sembach, K. R. 1996, *ARAA*, 31, 279
 Schneider, D. P., et al. 1993, *ApJS*, 87, 45
 Spitzer, L. 1978, *Physical Processes in the Interstellar Medium* (New York: John Wiley & Sons, Inc.)
 van Driel, W., van Woerden, H., Schwarz, U. J., and Gallagher, J. S., III. 1988, *A&A*, 191, 201
 Viegas, S. M. 1995, *MNRAS*, 276, 268